

WHAT YOU NEED TO KNOW ABOUT SCIATICA

The term sciatica describes the symptoms of leg pain and possibly tingling, numbness or weakness that originates in the lower back and travels through the buttock and down the large sciatic nerve in the back of the leg. Sciatica (pronounced sigh-at-ih-kah) is not a medical diagnosis in and of itself - it is a *symptom of an underlying medical condition*.

SCIATICA NERVE PAIN

Sciatica is often characterized by one or more of the following symptoms:

- Constant pain in only one side of the buttock or leg (rarely can occur in both legs)
- Pain that is worse when sitting
- Burning or tingling down the leg (vs. a dull ache)
- Weakness, numbness or difficulty moving the leg or foot
- A sharp pain that may make it difficult to stand up or to walk

Sciatic pain can vary from infrequent and irritating to constant and incapacitating. Specific sciatica symptoms also vary widely in type, location and severity, depending upon the condition causing the sciatica.

While symptoms can be very painful, it is rare that permanent sciatic nerve damage (tissue damage) will result.

THE SCIATIC NERVE AND SCIATICA

The sciatic nerve is the largest single nerve in the body and is composed of individual nerve roots that start by branching out from the spine in the lower back and combine to form the "sciatic nerve." When it is irritated, sciatica symptoms occur.

- The sciatic nerve starts in the lower back at lumbar segment 3 (L3).
- At each level of the lower spine a nerve root exits from the inside of the spine and then comes together to make up the large sciatic nerve.
- The sciatic nerve runs from the lower back, down the back of each leg
- Portions of the sciatic nerve then branch out in each leg to innervate certain parts of the leg - e.g. the buttock, thigh, calf, foot, toes.

The sciatica symptoms (e.g., leg pain, numbness, tingling, weakness, possibly symptoms that radiate into

Dr. Matthew Hartsburg

Address

Hartsburg Chiropractic

40 Lake Avenue Ext
Danbury, CT 06811

Telephone Number

203-792-4476

Qualifications

Graduated From: Life Chiropractic College
West

Years in Practice: 10

Treatments

- Chiropractic Manipulations
- Instrument Adjusting
- Decompression/Flexion Distraction
- Corrective Traction
- Physical Therapeutic Modalities
- Myofascial Release/Soft Tissue Therapy

"Why wait until you need to use risky drugs and surgery for your pain? When you understand the origin of the problem, permanent pain relief is possible naturally."

the foot) are different depending on where the nerve is pinched. For example, a lumbar segment 5 (L5) nerve impingement can cause weakness in extension of the big toe and potentially in the ankle.

THE COURSE OF SCIATICA PAIN

The incidence of sciatica increases in middle age. Rarely occurring before age 20, the probability of experiencing sciatic pain peaks in the 50s and then declines.

Often, a particular event or injury does not cause sciatica, but rather it tends to develop over time.

The vast majority of people who experience sciatica get better within a few weeks or months and find pain relief with non-surgical sciatica treatment. For others, however, sciatica pain from a pinched nerve can be severe and debilitating.

There are a few symptoms that may require immediate medical, and possibly surgical, intervention, such as progressive neurological symptoms (e.g. leg weakness) and/or bowel or bladder dysfunction (Cauda Equina Syndrome).

Because sciatica is caused by an underlying medical condition, treatment

is focused on relieving the underlying causes of symptoms. Treatment is usually self-care and/or non-surgical, but for severe or intractable cases surgery may be an option.

MOST COMMON CAUSES OF SCIATICA

- **Lumbar herniated disc** (also referred to as a slipped, ruptured, bulging, or protruding disc, or a pinched nerve) occurs when the soft inner core of the disc leaks out through the outer core and irritates the nerve root. Sciatica is the most common symptom of a lumbar herniated disc.
- **Degenerative disc disease** is diagnosed when a weakened disc results in excessive micro-motion at that spinal level, and inflammatory proteins from inside the disc become exposed and irritate the area (including the nerve roots).
- **Isthmic spondylolisthesis.** This condition occurs when a small stress fracture allows one vertebral body to slip forward on another (e.g. the L5 vertebra slips over the S1 vertebra).

The combination of disc space collapse, the fracture, and the

vertebral body slipping forward, can cause the nerve to get pinched and cause sciatica.

- **Lumbar spinal stenosis** is related to natural aging in the spine and is relatively common in adults over age 60. The condition typically results from a combination of one or more of the following: enlarged facet joints, overgrowth of soft tissue, and a bulging disc placing pressure on the nerve roots, causing sciatica pain.
- **Piriformis syndrome.** The sciatic nerve can get irritated as it runs under the piriformis muscle in the buttock. If the piriformis muscle irritates or pinches a nerve root that comprises the sciatic nerve, it can cause sciatica-type pain.
- **Sacroiliac joint dysfunction.** Irritation of the sacroiliac joint - located at the bottom of the spine - can also irritate the L5 nerve, which lies on top of the sacroiliac joint, causing sciatica-type pain.

Click here to read the full article:
<http://www.spine-health.com/conditions/sciatica/what-you-need-know-about-sciatica>

Talk to Dr. Hartsburg about specific questions related to your unique health situation.

Contact Dr. Hartsburg
to learn more or schedule an appointment
203-792-4476

<http://www.spine-health.com/doctor/chiropractor/matthew-hartsburg-danbury-ct>